

Schoolworkout Maths

Introduction to Probability: Lesson Notes and Examples

Key words-

Experiment

Outcome

Event

Example

Experiment: Pick a card from a standard pack of 52 cards

Outcomes: 2 of hearts, 9 of clubs, ace of spades, etc.

Event: picking a card with an even number on.

$$P(\text{even number}) = \frac{20}{52} = \frac{5}{13} \quad (2, 4, 6, 8, \text{ or } 10 \text{ of each suit})$$

Definition: An event is a set of possible outcomes.

Key results

Let A and B be two events.

✓ $0 \leq P(A) \leq 1$

Note: P(A) is short for the probability of A.

✓ $P(\text{A not happening}) = 1 - P(A)$

Note: A' means *not* A




✓ If A and B are *mutually exclusive* (so that they cannot occur at the same time), then

$$P(A \text{ or } B) = P(A) + P(B)$$

N.B. $P(A \cup B)$ means P(A or B)

✓ More generally, if A and B are not mutually exclusive:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$



Example: A university has 4000 first year undergraduates

2800 live in halls

1000 live in privately-rented accommodation

500 study medicine

340 of the medical students live in halls.

A student is picked at random.

Let H be the event the student lives in halls; R in privately-rented accommodation; M that they study medicine. Then the above information can be represented by the probabilities:

$$P(H) = \frac{2800}{4000} = \frac{7}{10}$$

$$P(R) = \frac{1000}{4000} = \frac{1}{4}$$

$$P(M) = \frac{500}{4000} = \frac{1}{8}$$

$$P(M \cap H) = \frac{340}{4000} = \frac{17}{200}$$

So, $P(H \cup R) = \frac{7}{10} + \frac{1}{4} = \frac{19}{20}$ (you can add the probabilities since H and R are mutually exclusive – you can't live in both types of accommodation at the same time).

But $P(H \cup M) = P(H) + P(M) - P(H \cap M)$
 $= \frac{7}{10} + \frac{1}{8} - \frac{17}{200} = \frac{37}{50}$ (H and M are not mutually exclusive – you can be a medical student and live in halls).

Independent events

A and B are *independent* if the probability of one happening isn't affected by whether the other happens or not. If A and B are independent, then:

$$P(A \cap B) = P(A) \times P(B).$$

Example: A dice is thrown twice. Find the probability

- of getting a 4 on both occasions;
- that neither number is a 2;
- both numbers are the same.

a) When you throw a dice twice, the outcomes of the throws are independent of each other. So

$$P(4 \text{ AND } 4) = \frac{1}{6} \times \frac{1}{6} = \frac{1}{36}$$

$$b) P(\text{not } 2 \text{ AND not } 2) = \frac{5}{6} \times \frac{5}{6} = \frac{25}{36}$$

$$c) P(\text{both numbers are the same}) = P(1, 1) + P(2, 2) + \dots + P(6, 6) = \frac{1}{36} + \frac{1}{36} + \dots + \frac{1}{36} = \frac{1}{6}$$

Some key techniques:

Sample space diagrams:

Eg two dice are each numbered 1, 2, 2, 3, 3, 6. They are both thrown and their scores are added together.

There are 36 equally likely combinations of scores:-

$$P(\text{total is 5}) = \frac{6}{36} = \frac{1}{6}$$

$$P(\text{total is even}) = \frac{18}{36} = \frac{1}{2}$$

$$P(\text{total score is at least 6}) = \frac{15}{36} = \frac{5}{12}$$

The events of winning a prize in the raffle and in the tombola are independent of each other.
Therefore:

$$\text{a) } P(2 \text{ prizes}) = P(R \text{ AND } T) = \frac{1}{80} \times \frac{1}{10} = \frac{1}{800}$$

$$\text{b) } P(\text{at least one prize}) = 1 - P(\text{no prizes}) = 1 - \frac{79}{80} \times \frac{9}{10} = \frac{89}{800}$$